

QUESTIONS FOR YOUR EYE DOCTOR

1. I answered Yes to either or both questions on the free, at-home Amsler test. What are my options for improving my vision?
2. I have had diabetes for ___ year(s). Will this affect the treatment types available to me?
3. I have high blood sugar. Will that increase the side effects of ILUVIEN?
4. I have had eye surgery before. How will this affect my treatment options?
5. How can I reduce the number of eye injections I currently receive?
6. What are the long term benefits of fewer eye injections?
7. What is the procedure like for ILUVIEN?
8. Can I still get ILUVIEN even though I have glaucoma, cataracts, or another eye condition?
9. What type of payment or insurance coverage is available?
10. What is the quality of life with DME? How does it progress?
11. Can DME be reversed?
12. Are there steps I can take to help slow the progression of DME?
13. How does my diabetes affect DME?

Indication

ILUVIEN® (fluocinolone acetonide intravitreal implant) 0.19 mg is indicated for the treatment of diabetic macular edema (DME) in patients who have been previously treated with a course of corticosteroids and did not have a clinically significant rise in intraocular pressure.

Important Safety Information

Do not use ILUVIEN if you have or think you might have an infection in or around the eye.

ILUVIEN should not be used if you have advanced glaucoma.

You should not use ILUVIEN if you are allergic to any ingredients of ILUVIEN.

Injections into the vitreous in the eye are associated with a serious eye infection (endophthalmitis), eye inflammation, increased eye pressure, glaucoma, and retinal detachments. Your eye doctor should monitor you regularly after the injection.

Use of corticosteroids including ILUVIEN may produce cataracts (ILUVIEN 82%; sham 50%), increased eye pressure (ILUVIEN 34%; sham 10%), glaucoma, and may increase secondary eye infections due to bacteria, fungi, or viruses. Let your doctor know if you have a history of herpes viral infections of the eye.

If the posterior capsule of the lens of your eye is missing or torn the ILUVIEN implant may move to the front chamber of the eye.

The most common side effects reported in patients with diabetic macular edema who were treated with ILUVIEN include cataracts (ILUVIEN 82%; sham 50%) and increased eye pressure (ILUVIEN 34%; sham 10%).

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.